

# Clinical Practice Competence and Its Associated Factors among Undergraduate Health Sciences Students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia, 2022

Telksew Yelma Yezengaw (✉ [yelmatelksew@gmail.com](mailto:yelmatelksew@gmail.com))

Harar Ethiopia

Simachew Animen

Bahir Dar University

Almaz Aklilu

Harar Ethiopia, Bahir Dar University

Wondu Feyisa

Harar Ethiopia, Bahir Dar University

---

## Research Article

**Keywords:** Competence, Clinical practice, Clinical practice competence

**Posted Date:** December 7th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-2351251/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Introduction:** Competence is the ability to perform a task with the desired result. Clinical experience is necessary to achieve the highest level of clinical competence after appropriate theoretical and practical training. It is estimated that many women and newborns die every year around the world due to a lack of qualified medical personnel. This is usually because women do not have access to competent health professionals.

**Objective:** This study aimed to assess clinical practice competence and its associated factors among undergraduate health sciences students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022.

**Methods:** Institutional based cross-sectional study was conducted from August 12-24/2022 among 403 undergraduate health sciences students at Bahir Dar Health Sciences Colleges. Simple random sampling technique was applied to select study subject and the data was collected by using a self-administered questionnaire. Data template format were prepared and entered into Epi Data version 4.1 and exported to SPSS version 25 for analysis. In binary logistic regression of dependent and independent variables with P-value,  $\leq 0.2$  were included in multi-variable logistic regression and  $\leq 0.05$  were considered as significant in this study.

**Result:** The prevalence of clinical practice competency among the study participant was 147(36.5%). students who were provide logbook [(AOR=5.40, 95% CI 2.91, 10.02)], adequate clinical cases in the clinical practice placement [(AOR=2.72, 95% CI 1.60, 4.60)], preceptor show different procedures [(AOR=2.50, 95% CI 1.33, 4.71)], student's confidence during conducting procedure [(AOR=4.16, 95% CI 1.67, 10.35)] and the suitability of the way of teaching to the learning styles of students during skills demonstration [(AOR=2.10, 95% CI 1.00, 4.40)] were significantly associated with clinical practice competence.

**Conclusion:** About 147 (36.5%) of study participants were clinically competent which was higher than previous studies conducted in Ethiopia. providing logbooks, adequate clinical cases, preceptor show different procedures, student's confidence and suitability of the way of teaching to the learning styles of students were significantly associated with clinical practice competence. Implementing logbooks, selection of clinical site, enhancing confidence of students, preferred teaching and learning styles and clinical preceptor support are important to improve the clinical competence of students.

## Introduction

The term competence has been derived from the Latin word "competent" that is to say ability and authorization. The concept of competence and its utilization within the nursing profession was mentioned by Banner (1984) in which nursing competence is the ability to perform a task with desirable outcomes (1).

Although they will sound similar, competence and competency don't seem to be necessarily synonymous. Competence may be a description of actions that will be demonstrated or observed and assessed, but competency refers to the skill itself (2).

Nursing competence has two approaches. The primary approach is Behaviorist, which focuses on skills, supports the direct observation of performance, and depends on the amount of each specific competence. Another approach is the Holistic approach, which regards competence in terms of a broad cluster of abilities that are conceptually linked and focuses on general attributes that are essential to effective performance (1).

The government of Ethiopia has developed different strategies to extend the competence of health professionals. The tutorial curriculum was changed from 3 years to a 4-year program and pre-service education has been started. But the supply of pre-service education might not translate into improved competence unless they master the essential knowledge and skills during their pre-service education (6-8).

The clinical learning environment could be a complex and dynamic place where the theoretical components of the curriculum are often integrated with the sensible and transformed into professional skills. It includes everything that surrounds students and affects their professional development within the clinical setting. Therefore, clinical practice occurs through interaction among students, educators, staff, patients, and environments (9, 10).

The result of effective learning in a health-care setting is achievement of clinical practice competency, which is the skill to successfully apply professional knowledge, attitudes and skills to new circumstances as well as familiar ones (12, 13)

Even though, the previous studies done in Ethiopia suggested that most new bachelor graduates have a lack of competence in the clinical environment (3, 6, 11). None of them have assessed whether simulation based learning factors have association with clinical competence or not, study will be to assess the clinical practice competence and associated factors among Midwifery and Nursing students at Bahir Car Health Sciences Colleges.

The results of the study will be useful in designing appropriate clinical practice implementation guidelines for nursing, midwifery education, the higher officials and nongovernmental organizations, the ministry of health, teaching institutions, and other stakeholders. Thus, a more supportive and relevant intervention can be implemented to help them achieve a higher level of clinical competence, contributing to society getting quality health care. It will also serve as a reference for future studies on the subject.

## **Methods**

### **Study design and period**

An institutional based cross-sectional study design was conducted from August 12 to 24/2022 at Bahir Dar Health Sciences Colleges.

### **Study setting**

The study was carried out in Bahir Dar. Bahir Dar is the capital city of the Amhara region in northern Ethiopia. It is a port on the south shore of the huge inland Lake Tana, and it has a population of around 332,856. The city is located approximately 490.34 km north-northwest of Addis Ababa (53).

The city has a number of public and private universities and colleges, including Bahir Dar University, Alkan University College, Bahir Dar Health Science College, GAMBY College of Medical Sciences, Kea-Med Medi College, Rift Valley University and Capital College of Business, and Health Science are the most (54).

### **Source Population**

All third and fourth year Bachelor of Science in Midwifery and Nursing students in Bahir Dar University and Bahir Dar health Sciences Colleges.

### **Study Population**

Randomly selected Bachelor of Science in Midwifery and Nursing Students

### **Inclusive and Exclusive Criteria**

#### **Inclusive Criteria**

All Third and Fourth year Bachelor of Science in Midwifery and Nursing students in Bahir Dar University and Bahir Dar health Sciences Colleges.

#### **Exclusive Criteria**

Third and Fourth year Bachelor of Science in Midwifery and Nursing students in Bahir Dar University and Bahir Dar health Sciences Colleges who withdraw or dropped out clinical placement during data collection period.

### **Sample Size Determination**

The sample size was computed based on the single population proportion calculation formula. The prevalence of clinical practice competence in the study done at Dilla University in 2016 was used to determine the sample size of 39.3% (6). The sample size for the second specific objective of this study to assess factors associated with clinical practice competence was determined by considering factors that are significantly associated with the outcome variable at ( $p < 0.05$ ), two-sided confidence level of 95%, margin of error of 5%, power of 80% and ratio of exposed to unexposed 1:1 using EPI Info Version 7. Finally, the required sample size for this particular study is decided by taking the maximum from the

calculated sample size for the first objective 366 plus 10% non-response rate (37), and then the final required sample size was 403.

## **Sampling Procedure**

Bahir Dar University and all Bahir Dar health Sciences Colleges of all Third and Fourth year Bachelor of Science in Midwifery and Nursing students were included in the study. The total sum of students in health sciences colleges' is 878, from those 433 are the 3rd and the remaining 445 are 4th year Midwifery and Nursing students. The total sample size was proportionally allocated to the university and each health Sciences College. The lists of students were obtained from the respective university and Health Sciences Colleges registrar. Then, the study participants from the university and each health Sciences College were selected by simple random sampling technique. The randomness of the selection was ensured until we get the required sample size for each department.

## **Study Variables**

### **Dependent variable:**

Clinical practice competence

### **Independent variables:**

- Socio-demographic factor (Age, Sex, Religion, Educational status of family, Residence and Year of study)
- Clinical instructors related factors
- Assessment method related factors
- Ward environment related factors
- Clinical staff related factors
- Students related factors
- Preceptor related factors
- Simulation-based learning related factors

## **Operational Definitions**

- Clinical practice competent: those students who scored mean and above the mean score of all competency domain assessment questions (6).
- Clinical practice incompetent: those students who scored below the mean score of all competency domain assessment questions (6).
- Clinical instructors: a person who provides direct supervision and instruction to students in the clinical aspect of health training education.

- Preceptors: an experienced licensed clinician who supervises health students during their clinical rotations, assign from institutions or colleges.
- Simulation-based learning: an educational activity that utilizes simulation aides to replicate clinical scenarios.
- Health sciences students: in this study refers to students who choose to pursue a degree in health science with Midwifery and Compressive Nurse.

## **Method of Data Collection**

The data were collected by a self-administered questionnaire prepared in English. Questionnaires are partially adapted from (3, 4, 6, 11, 25, 35, 40, 43). The data were collected by four B.Sc. in public health officers and supervised by two M.Sc. (a nurse and a midwife). The principal investigator from August 12 to 24/2022 supervised the data collection. Before the actual data collection, four data collectors were obtained one-day training about the aim of the study and the content of the instrument. Therefore, the data collectors were familiar with each question. It were considered as a mechanism for minimizing bias during the process of the data collection period.

## **Data Quality Assurance**

The data that has been collected is checked for completeness, accuracy, clarity and consistency after being processed neatly. 5% or 21 third and fourth year undergraduate courses in Midwifery and Nursing at Harar College of Health Sciences and Haramaya University are implemented, and the equipment is modified accordingly. Errors, ambiguities or incompleteness found are immediately corrected. Data collectors received a day's training on the content of the questionnaire, the method of data collection, and the purpose of the research. Supervisors and researchers monitor the data collection process during data collection. In addition, when you log into the computer, the data is checked before analysis. Reliability analysis was conducted to check the internal consistency of the 15 items of the competence measurement instrument, and the overall value of Cronbach's alpha was 0.915, indicating the very high consistency of the instrument to measure competence.

## **Data processing and Analysis**

Data were coded, cleaned and entered by Epi. Data was analyzed using computer database software version 4.1 and entered into statistical software SPSS version 25. Descriptive statistics such as frequency and percentage were used to represent dependent and independent variables. Frequency tables and graphs are used to present descriptive results. A bivariate logistic regression model was fitted as the most analytical method for this study. For the factors considered in this study, comparisons were made with 95% confidence intervals (CI) to predict clinical practice skills. Independent factors with a value of  $P < 0.2$  were included in the multivariate logistic regression model. Finally, the most significant associated factors were identified by multivariate logistic regression analysis. Then, adjusted odds ratios (AORs) with 95% confidence intervals were calculated for the most predictable variables, and statistical significance was accepted ( $P < 0.05$ ).

# Results

A total of 403 students have participated in the study, giving a response rate of 100%. The total, 213 (52.9%) of the respondents were females and the minimum and maximum age of the participants were 18 and 29 respectively. The largest numbers of students belonging to the age category of 20-24, 348 (86.4%) Half of the study participants 270 (67%) were Orthodox followers and 304 (75.4%) of them were Amhara by ethnicity (Table 1).

**Table 1: Socio-demographic characteristics of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Characteristics           |                                   | Frequency | Percentage % |
|---------------------------|-----------------------------------|-----------|--------------|
| University/College        | Bahir Dar University              | 69        | 17.1         |
|                           | Rift valley University            | 77        | 19.1         |
|                           | Alkan University College          | 69        | 17.1         |
|                           | GAMBY College of Medical Sciences | 118       | 29.3         |
|                           | Kea-Med Medical College           | 70        | 17.4         |
| Religion                  | Orthodox                          | 270       | 67.0         |
|                           | Muslim                            | 41        | 10.2         |
|                           | Protestant                        | 81        | 20.1         |
|                           | Others                            | 11        | 2.7          |
| Marital status            | Single                            | 392       | 97.3         |
|                           | Married                           | 11        | 2.7          |
| Ethnicity                 | Amhara                            | 304       | 75.4         |
|                           | Gurage                            | 29        | 7.2          |
|                           | Somalie                           | 16        | 4.0          |
|                           | Tigray                            | 4         | 1.0          |
|                           | SNNPR                             | 26        | 6.5          |
|                           | Oromo                             | 24        | 5.9          |
| Residence                 | Urban                             | 326       | 80.9         |
|                           | Rural                             | 77        | 19.1         |
| Departments               | Midwifery                         | 173       | 42.9         |
|                           | Nursing                           | 230       | 57.1         |
| Year of study             | 3 <sup>rd</sup> year              | 199       | 49.4         |
|                           | 4 <sup>th</sup> year              | 204       | 50.6         |
| Father educational status | Unable to write and read          | 36        | 8.9          |
|                           | Able to write and read            | 149       | 37.0         |
|                           | Primary (1-8)                     | 24        | 6.0          |
|                           | Secondary (9-12)                  | 47        | 11.7         |
|                           | Diploma and above                 | 147       | 36.5         |



|                                  |                          |     |      |
|----------------------------------|--------------------------|-----|------|
| <b>Mother educational status</b> | Unable to write and read | 72  | 17.9 |
|                                  | Able to write and read   | 159 | 39.5 |
|                                  | Primary (1-8)            | 35  | 8.7  |
|                                  | Secondary (9-12)         | 53  | 13.2 |
|                                  | Degree and above         | 84  | 20.8 |

### Prevalence of clinical practice competence

This study revealed that the overall clinical practice competence of the study participant was 147(36.5%) with 95% CI 32 - 41% (Figure 4).

### Clinical practice competencies by year of study and departments

There was a difference in clinical practice competence of the study participants in terms of year of study and departments. Regarding the department of study participants, 64 (43.5%) of Midwifery students and 83 (56.5%) of nursing students were clinically competent. Concerning year of study when increase in year the clinically competent level should be also, increase (Figure 2).

### Clinical instructor factors

Among the participant, more than half of them 348 (86.4%) were agreed on clinical instructor use different learning methods, and 128(31.8%) of them were disagreed regarding clinical instructor Orient the objective of clinical practice during clinical practice. Regarding the clinical instructor, 327(81.1%) of the participants were agreed on Spent enough time on clinical site and 117(29.0 %) of the participant were disagreed on clinical instructor provides constructive feedback (Table 2).

**Table 2: Clinical instructor characteristics response of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Clinical instructor factor |                                           | Yes       |      | No        |      |
|----------------------------|-------------------------------------------|-----------|------|-----------|------|
|                            |                                           | Frequency | %    | Frequency | %    |
| 1                          | Provide logbook                           | 285       | 70.7 | 118       | 29.3 |
| 2                          | Orient the objective of clinical practice | 275       | 68.2 | 128       | 31.8 |
| 3                          | Spent enough time on clinical site        | 327       | 81.1 | 76        | 18.9 |
| 4                          | Use different learning methods            | 348       | 86.4 | 55        | 13.6 |
| 5                          | Show clinical procedure                   | 323       | 80.1 | 80        | 19.9 |
| 6                          | Provides constructive feedback            | 286       | 71.0 | 117       | 29.0 |

### Clinical practice environment factors

Regarding the clinical practice environment is conducive, 306 (75.9%) of the study participant were agreed and 251(62.3%) of them were disagreed on clinical placement has a room for joint meetings during clinical practice. Concerning staff encourages students during clinical practice, 251(62.3%) of the study participant were agreed and 168 (41.7%) of them were disagreed on Clinical practice environment has sufficient cases (Table 3).

**Table 3: Clinical practice environment characteristics response of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Clinical practice environment factors |                                                                        | Yes       |      | No        |      |
|---------------------------------------|------------------------------------------------------------------------|-----------|------|-----------|------|
|                                       |                                                                        | Frequency | %    | Frequency | %    |
| 1                                     | Clinical practice environment is conducive                             | 306       | 75.9 | 97        | 24.1 |
| 2                                     | Clinical practice environment has sufficient cases                     | 235       | 58.3 | 168       | 41.7 |
| 3                                     | Clinical practice environment has sufficient material                  | 237       | 58.8 | 166       | 41.2 |
| 4                                     | Clinical practice environment has meet objectives of clinical practice | 251       | 62.3 | 152       | 37.7 |
| 5                                     | Clinical practice environment has sufficient wards                     | 246       | 61.0 | 157       | 39.0 |
| 6                                     | Clinical placement has a room for joint meetings                       | 162       | 40.2 | 241       | 59.8 |
| 7                                     | Staff allows students to perform tasks during clinical practice        | 249       | 61.8 | 154       | 38.2 |
| 8                                     | Staff encourages students during clinical practice                     | 251       | 62.3 | 152       | 37.7 |

## Assessment method factors

Regarding assessment methods has a positive influence on clinical practice, 289(71.7%) of participants were agreed and 245(60.8%) of them were disagreed as assessment methods address the three learning domains. About 282 (70.0%) of the participants were agreed on instructor orient about assessment methods during clinical practice and 199(49.4%) of them were disagreed on instructor use checklist to assess the performance during clinical practice (Table 4).

**Table 4: Assessment method characteristics of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Assessment method factors |                                                                 | Yes       |      | No        |      |
|---------------------------|-----------------------------------------------------------------|-----------|------|-----------|------|
|                           |                                                                 | Frequency | %    | Frequency | %    |
| 1                         | Instructors orient about assessment                             | 282       | 70.0 | 121       | 30.0 |
| 2                         | Assessment method has a positive influence on clinical practice | 289       | 71.7 | 114       | 28.3 |
| 3                         | Instructor uses continuous assessment                           | 224       | 55.6 | 179       | 44.4 |
| 4                         | Assessment methods address the three learning domain            | 158       | 39.2 | 245       | 60.8 |
| 5                         | Instructor use checklist to assess                              | 204       | 50.6 | 199       | 49.4 |

## Preceptor factors

Majority 391(97.0%) of the respondents agreed that have preceptor in your clinical practice and 122 (31.2%) of the participant have disagreed regarding preceptor provides constructive feedback during clinical practice (Table 5).

**Table 5: Preceptor factor characteristics of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Preceptor factor                                  | Yes       |      | No        |      |
|---------------------------------------------------|-----------|------|-----------|------|
|                                                   | Frequency | %    | Frequency | %    |
| 1 Do you have preceptor in your clinical practice | 391       | 97.0 | 12        | 3.0  |
| 2 Preceptor identifies student's learning needs   | 357       | 91.3 | 34        | 8.6  |
| 3 Preceptor has good clinical skills              | 322       | 82.3 | 69        | 17.6 |
| 4 Preceptor show different procedures             | 309       | 79.0 | 82        | 20.9 |
| 5 Preceptor provides constructive feedback        | 269       | 68.7 | 122       | 31.2 |
| 6 Preceptor has good interpersonal skills         | 270       | 69.0 | 121       | 30.9 |

### Student factors

Regarding the absence during clinical practice 321(79.7%) were absent from clinical practice without permission, out of them 281(87.5%) were absent once, 34(10.5%) were twice and 6 (1.8%) were three times and above (Table 6).

**Table 6: Student factor characteristics of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Student factor during clinical practice                    | Yes       |       | No        |      |
|------------------------------------------------------------|-----------|-------|-----------|------|
|                                                            | Frequency | %     | Frequency | %    |
| 1 Having a good relationship with team members             | 335       | 83.1  | 68        | 16.9 |
| 2 Having good information exchange habit with team members | 336       | 83.4  | 67        | 16.6 |
| 3 I was motivated and eager to learn                       | 403       | 100.0 | -         | -    |
| 4 I was confident during conducting procedure              | 356       | 88.3  | 47        | 11.7 |
| 5 Absent from clinical practice without permission         | 321       | 79.7  | 82        | 20.3 |
| 6 I was punctual during my clinical practice               | 365       | 90.6  | 38        | 9.4  |
| 7 My parent economic status affected my clinical practice  | 168       | 41.7  | 235       | 58.3 |

### Simulation-based learning factors

Majority 340(84.4%) of the respondents agreed that get necessary help in the use of equipment during skills demonstration and 162 (40.2%) of the participant have disagreed regarding programs of skills demonstration are flexible and adjustable for simulation class (Table 7).

**Table 7: Simulation-based factor characteristics of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Simulation-based learning related factor |                                                                                             | Yes       |      | No        |      |
|------------------------------------------|---------------------------------------------------------------------------------------------|-----------|------|-----------|------|
|                                          |                                                                                             | Frequency | %    | Frequency | %    |
| 1                                        | Teacher provides me accurate information about skills requirements                          | 324       | 80.4 | 79        | 19.6 |
| 2                                        | Teacher explains learning objectives for simulation-learning at the beginning of the period | 286       | 71.0 | 117       | 29.0 |
| 3                                        | Teacher gives me enough time to meet the objective(s)                                       | 290       | 72.0 | 113       | 28.0 |
| 4                                        | Teacher assists me in developing long-term skills                                           | 286       | 71.0 | 117       | 29.0 |
| 5                                        | Students' different backgrounds are taken into account                                      | 274       | 68.0 | 129       | 32.0 |
| 6                                        | The way my teachers taught the simulation is suitable to the way I learn                    | 336       | 83.4 | 67        | 16.6 |
| 7                                        | The teaching methods used in the simulation are helpful and effective                       | 306       | 75.9 | 97        | 24.1 |
| 8                                        | There are enough skills practicing programs per semester                                    | 248       | 61.5 | 155       | 38.5 |
| 9                                        | Programs of skills demonstration are flexible and adjustable for simulation class           | 241       | 59.8 | 162       | 40.2 |
| 10                                       | Number of students per teaching group is small                                              | 248       | 61.5 | 155       | 38.5 |
| 11                                       | I can get necessary help in the use of equipment                                            | 340       | 84.4 | 63        | 15.6 |
| 12                                       | I can assess my own skills performance critically                                           | 312       | 77.4 | 91        | 22.6 |
| 13                                       | My teacher gives me necessary feedback                                                      | 337       | 83.6 | 66        | 16.4 |

### Factors associated with clinical practice competence

Hosmer and lemeshow model of fitness show that the model was adequately fit with p-value of 0.429 which is greater than 0.05.

In bivariate logistic regression analysis, provide logbook, show clinical procedure, clinical practice environment has sufficient cases, staff encourages students during clinical practice, assessment method has a positive influence on clinical practice, instructor use checklist to assess the performance, preceptor show different procedures, confident during conducting procedure, way of teachers taught the simulation is suitable to the way of learn, number of students per teaching group is small and get necessary help in the use of equipment were found to be associated factors with clinical practice competence in bivariate with p-value less than 0.2. Thus, transferred to multivariable logistic regression model. In multivariable model provide logbook, clinical practice environment has sufficient cases, assessment method has a positive influence on clinical practice, preceptor show different procedures, confident during conducting procedure and the way of teachers taught the simulation is suitable to the way of learn where maintain their association with clinical practice competence.

Students who were providing logbook were 5 times more likely competent than those who were not logbooks [(AOR=5.40, 95% CI 2.91, 10.02)].

This study also revealed that students with adequate clinical cases in the clinical practice placement were 2 times clinically competent than those students with inadequate clinical cases in the clinical practice placement with [(AOR=2.72, 95% CI 1.60, 4.60)].

Preceptor show different procedures was significantly associated with the clinical practice competence of students. Students who were different procedures showed by preceptors during clinical practice were 2 times more likely competent than those who have not shown different procedures [(AOR=2.50, 95% CI 1.33, 4.71)].

Among student factors included in the study, student's confidence during conducting procedure was significantly associated with the clinical practice competency of students. Students who were confident during conducting a procedure were 4 times more likely competent than those who were not confident during conducting a procedure [(AOR=4.16, 95% CI 1.67, 10.35)].

The way of teachers taught the simulation is suitable to the way of learn was significantly associated with the clinical practice competence of students. During skills demonstration, the way of teachers taught the simulation is suitable to the way to learn were 2 times more likely competent than those large number of students per teaching group [(AOR=2.10, 95% CI 1.00, 4.40)].

**Table 8: Results of bivariate and multivariate analysis for clinical practice competence and associated factors among Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).**

| Variables                                                       |     | Clinical practice competence |                      |                      |                         |
|-----------------------------------------------------------------|-----|------------------------------|----------------------|----------------------|-------------------------|
|                                                                 |     | Clinically Incompetent       | Clinically Competent | COR (95%CI)          | AOR (95%CI)             |
| Instructor provide clinical practice logbook                    | Yes | 166<br>(41.2%)               | 119<br>(29.5%)       | 2.30<br>(1.41, 3.74) | 5.40 (2.91, 10.02) ***  |
|                                                                 | No  | 90<br>(22.3%)                | 28<br>(6.9%)         | 1                    | 1                       |
| Instructor show clinical procedure                              | Yes | 197<br>(48.9%)               | 126<br>(31.3%)       | 1.79<br>(1.04, 3.10) | 1.63 (0.71, 3.74)       |
|                                                                 | No  | 59<br>(14.6%)                | 21<br>(5.2%)         | 1                    | 1                       |
| Clinical placement has sufficient cases                         | Yes | 157<br>(39.0%)               | 78<br>(19.4%)        | 0.71<br>(0.47, 1.07) | 2.72(1.60, 4.60) ***    |
|                                                                 | No  | 99<br>(24.6%)                | 69<br>(17.1%)        | 1                    | 1                       |
| Staff encourages students during clinical practice              | Yes | 170<br>(42.2%)               | 81<br>(20.1%)        | 0.62<br>(0.41, 0.94) | 1.40 (0.75, 2.61)       |
|                                                                 | No  | 86<br>(21.3%)                | 66<br>(16.4%)        | 1                    | 1                       |
| Assessment method has a positive influence on clinical practice | Yes | 194<br>(48.1%)               | 95<br>(23.6%)        | 0.58<br>(0.37, 0.90) | 2.147(1.251, 3.684) *** |
|                                                                 |     | 62                           | 52                   |                      |                         |

|                                                                           |     |                |                |                      |                         |
|---------------------------------------------------------------------------|-----|----------------|----------------|----------------------|-------------------------|
|                                                                           | No  | (15.4%)        | (12.9%)        | 1                    | 1                       |
| Instructors use checklist to assess performance                           | Yes | 131<br>(32.5%) | 73<br>(18.1%)  | 0.51<br>(1.00, 2.29) | 0.71 (0.40, 1.27)       |
|                                                                           | No  | 125<br>(31.0%) | 74<br>(18.4%)  | 1                    | 1                       |
| Preceptor show different procedures                                       | Yes | 197<br>(48.9%) | 124<br>(30.8%) | 1.61<br>(0.94, 2.74) | 2.50<br>(1.33, 4.71)**  |
|                                                                           | No  | 59<br>(14.6%)  | 23<br>(5.7%)   | 1                    | 1                       |
| Confident during performing procedure                                     | Yes | 220<br>(54.6%) | 136<br>(33.7%) | 2.02<br>(0.99, 4.10) | 4.16<br>(1.67, 10.35)** |
|                                                                           | No  | 36<br>(8.9%)   | 11<br>(2.7%)   | 1                    | 1                       |
| The way of teachers taught the simulation is suitable to the way of learn | Yes | 211<br>(52.4%) | 125<br>(31.0%) | 1.21<br>(0.69, 2.11) | 2.10 (1.00, 4.40)*      |
|                                                                           | No  | 45<br>(11.2%)  | 22<br>(5.5%)   | 1                    | 1                       |
| Number of students per teaching group is small                            | Yes | 148<br>(36.7%) | 100<br>(24.8%) | 0.64<br>(0.42, 0.98) | 1.10 (0.64, 1.91)       |
|                                                                           | No  | 108<br>(26.8%) | 47<br>(11.7%)  | 1                    | 1                       |
| During skills demonstration, I can get                                    |     | 216            | 124            |                      |                         |



|                                        |     |         |         |                      |                   |
|----------------------------------------|-----|---------|---------|----------------------|-------------------|
| necessary help in the use of equipment | Yes | (53.6%) | (30.8%) | 1.00<br>(0.57, 1.75) | 0.89 (0.43, 1.84) |
|                                        |     | 40      | 23      |                      |                   |
|                                        | No  | (9.9%)  | (5.7%)  | 1                    | 1                 |

P<0.05\*, P<0.01\*\*, P<0.001\*\*\*, COR= Crude Odd Ratio, AOD=Adjusted Odd Ratio, CI=confidence Interval, P-value set <0.2 for bivariate logistic regression and 0.05 for multivariate logistic regression

## Discussion

This study revealed that the overall clinical practice competence of the study participant was 147(36.5%) with 95% CI (32.0, 41.0%).

The result of this study was shows that lower than compared with the finding 65% in Bahir Dar University College of medicine and health science students in 2019. This discrepancy might be due to the difference in which the previous study was conducted in a single public institution, including all undergraduate students, but this study covers five public, and private institutions, only graduating midwifery and nursing students were included (26). In addition, this study result it is lower than compared with the finding 78.6% study done at Debre Birhan Health Science College in 2014. This discrepancy might be due to the difference in study setting and characteristics of study participants (23).

The result of this study is also inconsistent with a study conducted in Finland, which was 66.7%. This discrepancy may be due to the difference in sample size in which the previous study use small sample size compared to this study, socio-economic status of study participant and difference in curriculum of the nation (22).

The present study revealed that prevalence of clinical practice competence was almost similar as compared with the finding in two studies of 33.6% study done among graduating nursing students attending at Amhara Region Universities and 39.3% at Dilla University in 2016 (3, 6).

The result of this study shows that the prevalence of clinical practice competence were 36.5%. It is higher than the findings in two recent studies in Mettu University and Dire Dawa Health Sciences Colleges. This discrepancy might be due to the difference of study setting and characteristics of study participants (24, 25). In addition, this study result it is higher than compared with the finding of 23.2% among nursing and 22.8% among Midwifery student conducted in Hawassa University. This discrepancy might be due to the difference in which the previous study was conducted in a single public institution, including all undergraduate students, but this study covers five public, and private institutions, only graduating midwifery and nursing students were included (11).

In this study, students with adequate clinical case in practice placement has increase clinical practice competence (AOR=3.239, 95% CI 1.717, 6.110) than students with few clinical cases in practice placements. This is in line with the study done in Hawassa University in 2016 and Bahir Dar University College of medicine and health science students in 2019 (11, 26).

The above finding was inconsistent with a cross-sectional study done in Dilla University that was not significantly associated with the clinical practice competence of students. This variation might be due to the difference in characteristics of the respondent, sample size and study setting. The previous study was conducted in one governmental institution, the study participants were all health science students, but the current study was conducted in a private and governmental institution, and the study participants were Midwifery and Nursing students, and also it is the sample size was relatively smaller than the current study (6).

In this study, instructor provides clinical practice logbook during clinical practice are statistically significant factors. Those students who have logbook were five times more likely they are clinically competent compared with those who do not have logbook [(AOR=5.40, 95% CI 2.91, 10.02)]. This finding is supported by a study conducted in Assiut University and Shiraz University. This may be due to the reason that logbook effects on clinical learning (55, 56).

However, the above was inconsistent with a study done in Mettu University and Dire Dawa Health Sciences Colleges that was not significantly associated with the clinical practice competence of students. This discrepancy might be due to the difference of study setting and characteristics of study participants and in which might be due to lack of education on logbook in part of clinical teachers. Further study of the whole of the logbook would be useful (24, 25).

Preceptor show different procedures was significantly associated with the clinical practice competence of students. Students who were different procedures showed by preceptors during clinical practice were 2 times more likely competent than those who have not shown different procedures [(AOR=2.50, 95% CI 1.33, 4.71)]. This finding is in line with a study conducted in Amhara Region Universities, in Ghana, Indonesia and Pakistan (3, 37, 41, 42).

The above finding was inconsistent with a cross-sectional study done in Dilla University and Mettu University that was not significantly associated with the clinical practice competence of students. This variation might be due to the difference in characteristics of the respondent and study setting. The previous study was conducted in one governmental institution, the study participants were all health science students, but the current study was conducted in a private and governmental institution, and the study participants were Midwifery and Nursing students, and also they have preceptors in clinical practice.

Among student factors included in the study, student's confidence during conducting procedure was significantly associated with the clinical practice competency of students. Students who were confident during conducting a procedure were 4 times more likely competent than those who were not confident

during conducting a procedure [(AOR=4.16, 95% CI 1.67, 10.35)]. This finding is in line with a study conducted in Dire Dawa Health Sciences Colleges, Indonesia and Sweden. This may be due to the reason that self-confidence is a key component of effective clinical competence (25, 37, 38).

The above finding was inconsistent with a cross-sectional study done in Tanzania, in which the confidence of nursing students was not significantly associated with clinical practice competence. This variation might be due to the difference in sample size and the characteristics of the study participants, in which the sample size was 208 and the study participants were Diploma Nursing Students (35).

In the current study, the suitability of the way of teaching to the learning styles of students was significantly associated with the clinical practice competence of students. During skills demonstration, suitability of the way of teaching to the learning styles of students were 2 times more likely competent than those large number of students per teaching group [(AOR=2.10, 95% CI 1.00, 4.40)]. This finding is supported by a study conducted in public universities and colleges in Harar and Dire Dawa cities, Eötvös Loránd University, Indonesia, SMS & R, Sharda University, India and northwestern South Carolina (43, 55-58)

The above finding was inconsistent with a cross-sectional study done in Gondar University in which was not significantly associated. This variation might be due to the difference in characteristics of the respondent, sample size and study setting. The previous study was conducted the study participants were only midwifery students, but the current study was conducted in a study participants were Midwifery and Nursing students and also it is the sample size was smaller than the current study (59).

### **Research Limitation**

This study was includes both private and governmental institution compares to study done in northern Ethiopia. The study was also based on self-report by students and was not supported by qualitative methods.

## **Conclusions**

Generally, the results of this study revealed that about 147 (36.5%) of study participants were clinically competent, which was higher than previous studies conducted in Ethiopia. Providing logbooks during clinical practice, adequate clinical placement cases, preceptor show different procedures during clinical practice; student's confidence during conducting procedure during clinical and suitability of the way of teaching to the learning styles of students during skills demonstration were significantly associated with clinical practice competence of study participants.

## **Recommendations**

**Federal Ministry of Health (FMOH) and Ministry of Education**

The federal ministry of health and ministry of education should be training of core faculty in simulation-based teaching and assessment methods, and acquisition of a wide range of simulation equipment. Incorporating simulation based teaching activities designed to improve the skills, attitudes, and knowledge of the graduate students. In addition to that, the teaching - learning process are understanding the nature of individual difference and use the right strategies and method together with adopting teaching style of teachers, that it to get the smooth interaction in the classroom. This helps learners to develop their critical thinking, problem- solving and applying for their real life. Should work closely with a teaching institution and health facilities, other stakeholders to overcome those gaps.

### **To Bahir Dar University and Bahir Dar Health Science Colleges**

- The Colleges also give much infuses to quality of education by monitoring the flow of course and practical session.
- The Colleges must assign those clinical instructors and preceptors who have clinical experience.
- The Colleges must Strength the skill's lab (demonstration) room, and it should be together with focusing on the way of teaching of the teacher in terms of his or her teaching styles, the use of the method, technique, and strategy.
- The Colleges should take different actions to improve the student's confidence while they are conducting procedure and establish a team to overcome those gaps.
- The Colleges monitor before attaching the student's appropriate selection of clinical placement site.
- The Colleges develop a system to control instructors who miss practice schedule and integrating the clinical staff to academic staff
- The College's Implementation of logbook in clinical education for all clinical courses and inserts logbook as a method of evaluation of internships students during an internship training course in the midwifery and nursing bachelor program.

## **Abbreviations**

AOR, Adjusted Odds Ratio, BSc N/Mw, Bachelor of Science in Nursing/Midwifery, CI, Conference Interval, FMOH, Federal Ministry of Health, OR, Odds Ratio, SBL ,Simulation Based Learning, SPSS, Statistical Package for the Social Sciences, SSA, Sub Saharan Africa, WHO, World Health Organization

## **Declarations**

### **Data Sharing Statement**

All the data of this study are available from the corresponding author upon request.

### **Ethical Consideration**

Ethical approval was obtained from the Institutional Review Board of Bahir Dar University College of Medicine and Health Science with protocol number 517/2022. Formal letters were taken from the College of Medicine and Health Sciences and the Department of Midwifery, and letters are sent to the Colleges of Health Sciences and other relevant bodies for cooperation. The purpose of the study were explained to the study participants. During data collection, verbal consent it was obtained from participants to confirm their willingness to participate. Participants were informed that participation was voluntary and that they could withdraw at any time if they are not comfortable with the questionnaire. Names or personal identifiers will not be included in written questionnaires to ensure participant confidentiality.

## **Acknowledgment**

We are very grateful to Bahir Dar University for the financial support to this study and the department of midwifery for its mentorship. All study participants for their willingness in responding to our questionnaire.

## **Author Contributions**

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted and agree to be accountable for all aspects of the work.

## **Funding**

This work has been funded by Bahir Dar University for data collection purposes as for MSc thesis. College of Medicine and Health Sciences and the Department of Midwifery was involved in the project through monitoring and evaluation of the work from the beginning to the result submission. But this organization did not involve in designing, analysis, critical review of its intellectual content, preparation of the manuscript and the budget funded by this organization did not include for publication.

## **References**

1. Lejonqvist G-B. Clinical Competence: the Core of Nursing Education. 2018.
2. Schroeter K. Competence Literature Review, Competency & Credentialing Institute. October Denver. 2008.
3. Getie A, Tsige Y, Birhanie E, Tlaye KG, Demis A. Clinical practice competencies and associated factors among graduating nursing students attending at universities in Northern Ethiopia: institution-based cross-sectional study. *BMJ open*. 2021;11(4):e044119.
4. Ireland NaMBo. competence-assessment-tool-for-Nurses 2015. p. 27.

5. Fullerton JT, Thompson JB, Johnson P. Competency-based education: the essential basis of pre-service education for the professional midwifery workforce. *Midwifery*. 2013;29(10):1129-36.
6. Tesfaye TS, Alemu W, Mekonen T. Perceived clinical practice competency and associated factors among undergraduate students of medicine and health science collage in Dilla University, SNNPR, Ethiopia. *Advances in Medical Education and Practice*. 2020;11:131.
7. Yigzaw T, Ayalew F, Kim Y-M, Gelagay M, Dejene D, Gibson H, et al. How well does pre-service education prepare midwives for practice: competence assessment of midwifery students at the point of graduation in Ethiopia. *BMC medical education*. 2015;15(1):1-10.
8. Ahmed NG, Adam SM, Abd Al-Moniem II. Patient safety: assessing nurses' perception and developing an improvement plan. *Life Science Journal-Acta Zhengzhou University Overseas Edition*. 2011;8(2):53-64.
9. Papastavrou E, Dimitriadou M, Tsangari H, Andreou C. Nursing students' satisfaction of the clinical learning environment: a research study. *BMC nursing*. 2016;15(1):1-10.
10. Steven A, Magnusson C, Smith P, Pearson PH. Patient safety in nursing education: contexts, tensions and feeling safe to learn. *Nurse education today*. 2014;34(2):277-84.
11. Fikre R. Assessment of factors affecting clinical practice competency of undergraduate health science students in Hawassa University, South, Ethiopia. *Annals of Clinical and Laboratory Research*. 2016;4(1):0-.
12. Wong BSH. Clinical Competency: experience of new graduated nurses from bachelor degree of nursing in University Malaysia Sarawak. 2013.
13. Kylmänen P, Spasic A. Assessing competence in technical skills of theatre nurses in India and Sweden: Evaluation of an observational tool. 2010.
14. White AH. Clinical decision making among fourth-year nursing students: An interpretive study. *Slack Incorporated Thorofare, NJ*; 2003. p. 113-20.
15. Susan Sportsman R, editor *Competency education and validation in the United States: what should nurses know?* Nursing forum; 2010: Blackwell Publishing Ltd.
16. UNFPA-ICM. Investing in Midwives and others with Midwifery skills to save the lives of mothers and newborns and improve their health. 2006:9-13.
17. Carr C FV, Fogstad H, Garden B, Johnson P, Laski L, et al. . The state of the world's Midwifery delivering saving health lives. 2011:5-7.
18. Goshu M, Godefay H, Bihonegn F, Ayalew F, Haileselassie D, Kebede A, et al. Assessing the competence of midwives to provide care during labor, childbirth and the immediate postpartum period–A cross sectional study in Tigray region, Ethiopia. *PloS one*. 2018;13(10):e0206414.
19. Kasine Y, Babenko-Mould Y, Regan S. Translating continuing professional development education to nursing practice in Rwanda: Enhancing maternal and newborn health. *International journal of Africa nursing sciences*. 2018;8:75-81.
20. Ethiopia FDRo. Ethiopia mini demographic and health survey key indicators. 2019:13-5.

21. Feysia B, Herbst C, Lemma W. The health workforce in Ethiopia: addressing the remaining challenges. 2012.
22. Kajander-Unkuri S. Nurse competence of graduating nursing students. 2015.
23. Hailu A, Ditta H, Zewdie Z. Competency assessment and factors associated with it among health professionals at Debre Birhan Health Science College. *Open Journal of Nursing*. 2014;2014.
24. Amsalu B, Fekadu T, Mengesha A, Bayana E. Clinical practice competence of mettu university nursing students: a cross-sectional study. *Advances in Medical Education and Practice*. 2020;11:791.
25. Hailu M, Welday M, Haftu A, Tadesse D, Weldeamanel T, Amsalu B, et al. Clinical Practice Competence and its Associated Factors Among Midwifery and Nursing Students at Dire Dawa Health Sciences Colleges, East Ethiopia, 2020. *Advances in Medical Education and Practice*. 2021;12:1539.
26. Ayenew A. Clinical practice competency and associated Factors of undergraduate health science students in Bahir Dar University, Bahir Dar, Ethiopia. *Journal of Vaccines and Clinical Trials*. 2019; 4( 3 ):12-3.
27. Hakimzadeh R, Ghodrati A, Karamdost N, Ghodrati H, Mirmosavi J. Factors affecting the teaching-learning in nursing education. *GSE Journal of Education*. 2013;2013:174-84.
28. Bifftu BB, Dachew BA, Tadesse Tiruneh B, Mekonnen Kelkay M, Bayu NH. Perceived clinical competence among undergraduate nursing students in the university of Gondar and Bahir Dar university, Northwest Ethiopia: A cross-sectional institution based study. *Advances in Nursing*. 2016;2016.
29. Angasu K, Bekela T. Achievement of Clinical Learning Outcomes and Associated Factors Among Midwifery and Nursing Undergraduate Students of Jimma University, Ethiopia. *Advances in Medical Education and Practice*. 2021;12:987.
30. Amro NR, Al Sous A, Shkherat S, Nahli H, Hassasneh R, Slimi A. The theory Practices Gap among Nursing and Midwifery Students in Palestine. *International Journal of Innovative Research in Medical Science (IJIRMS)*. 2017;2(11).
31. Parvin N AF, Vardanjani LR, Dadkhah NK, Jouybari LM. Nurses attitude towards attendance of Nursing students in the clinical setting in Shahrekord. *Journal of Nursing and Midwifery Sciences*. 2016;3(4):22-4.
32. L S. Midwifery student experience at clinical site at univesrty of Sovenga. South Africa. 2013:24-6.
33. Gulbu Tanriverdi P, Katar T. Problems Experienced by Midwifery And Nursing Students in Turkey During Clinical Practice and their Recommended Solutions to the Problems. *International Journal of Caring Sciences*. 2009;2(1):22.
34. FAW G. Nursing and Midwifery Students Lived Experiences During Clinical Practice in Palestine. *International Journal of Scientific and Research*. 2017;2(7):45–7.
35. Gemuhay HM, Kalolo A, Mirisho R, Chipwaza B, Nyangena E. Factors affecting performance in clinical practice among preservice diploma nursing students in Northern Tanzania. *Nursing Research and Practice*. 2019;2019.

36. Dorothy A-P, Linda AS, Florence A. The attitudes of student nurses toward clinical work. *International Journal of Nursing and Midwifery*. 2013;5(2):22-7.
37. Rizany I, Hariyati RTS, Handayani H. Factors that affect the development of nurses' competencies: a systematic review. *Enfermeria clinica*. 2018;28:154-7.
38. Bäck L, Hildingsson I, Sjöqvist C, Karlström A. Developing competence and confidence in midwifery-focus groups with Swedish midwives. *Women and Birth*. 2017;30(1):e32-e8.
39. Aragaw Y, Sinishaw W, Daba W, Mekie M. Attitude of Nursing and Midwifery students towards clinical practice and its associated factors in Northwest Ethiopia: a cross-sectional study. *BMC research notes*. 2019;12(1):1-6.
40. Serrano-gallardo P M-mM, Espejo-matorrales F. . Factors associated to clinical learning in nursing students in primary health care. *Revista Lat americana Enferm*. 2016(24):5-6.
41. Rani S HM, Afzal M, Gillani SA. The Influence of personal characteristics of preceptor on professional grooming of Nursing students. *International Journal of Medical Research & Health Science*. 2019;8(5):91–3.
42. Atakro CA GJ. Preceptorship versus clinical teaching partnership. *Hindawi*. 2016(2016):2-4.
43. Jamie AH, Mohammed AA. Satisfaction with simulation-based education among Bachelor of Midwifery students in public universities and colleges in Harar and Dire Dawa cities, Ethiopia. *European journal of midwifery*. 2019;3:19.
44. Eldarir SA, A. Nagwa, and A. Hamid. . "Objective structured clinical evaluation (OSCE) versus traditional clinical students achievement at maternity nursing: A comparative approach.". *IOSR Journal of Dental and Medical Sciences*. 2013; 4.3:63-8.
45. Aimée MM. Life science journal-acta zhengzhou university overseas rwandan midwifery students benefits and barriers in the use of simulation based learning regarding neonatal resuscitation edition. *rwanda: university of rwanda*; 2019.
46. Asegid A, Assefa N. Effect of simulation-based teaching on nursing skill performance: a systematic review and meta-analysis. *Frontiers of Nursing*. 2021;8(3):193-208.
47. Ricketts B. The role of simulation for learning within pre-registration nursing education – A literature review. *Nurse Education Today*. 2011;31(7):650-4.
48. Shin S, Park J-H, Kim J-H. Effectiveness of patient simulation in nursing education: Meta-analysis. *Nurse Education Today*. 2015;35(1):176-82.
49. Hope A, Garside J, Prescott S. Rethinking theory and practice: Pre-registration student nurses experiences of simulation teaching and learning in the acquisition of clinical skills in preparation for practice. *Nurse Education Today*. 2011;31(7):711-5.
50. LISKI SA, O'DELL V. Integration of Theory and Practice: Experiential Learning Theory and Nursing Education. *Nursing Education Perspectives*. 2010;31(2):106-8.
51. Tilden VP, Tilden S. Benner, P. (1984). From novice to expert, excellence and power in clinical nursing practice. Menlo Park, CA: Addison-Wesley Publishing Company, 307 pp., \$12.95 (soft cover).



Research in Nursing & Health. 1985;8(1):95-7.

52. Nickless LJ. The use of simulation to address the acute care skills deficit in pre-registration nursing students: A clinical skill perspective. *Nurse Education in Practice*. 2011;11(3):199-205.
53. "Homepage of Lake Tana Biosphere Reserve" 2021 [updated 2 July 2021].
54. Bureau ARH. Amhara Regional Health Bureau. 2017.
55. Phavadee S. The way of teaching toward different students' learning styles/Teaching that takes into account different learning styles. *Opus et Educatio*. 2020;7(2).
56. Ridwan H, Sutresna I, Haryeti P, editors. Teaching styles of the teachers and learning styles of the students. *Journal of Physics: Conference Series*; 2019: IOP Publishing.
57. Kharb P, Samanta PP, Jindal M, Singh V. The learning styles and the preferred teaching–learning strategies of first year medical students. *Journal of clinical and diagnostic research: JCDR*. 2013;7(6):1089.
58. Wilson ML. Students' learning style preferences and teachers' instructional strategies: Correlations between matched styles and academic achievement: Liberty University; 2011.
59. Temesgen Worku Gudayu MBB, and Mengstu Melkamu Asaye. Self-Efficacy, Learner Satisfaction, and Associated Factors of Simulation Based Education among Midwifery Students: A Cross-Sectional Study. *Hindawi Publishing Corporation Education Research International*. 2015;2015:7.

## Figures

### OVERALL CLINICAL COMPETENCE




Figure 1

The overall clinical competence of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).




Figure 2

Clinical practice competencies by year of study of Midwifery and Nursing students at Bahir Dar Health Sciences Colleges, Northwest Ethiopia 2022, (n=403).